

RUMINAL BIOHYDROGENATION PATTERN OF POLY-UNSATURATED FATTY ACID AS INFLUENCED BY DIETARY TANNIN

Anuraga Jayanegara

*Department of Nutrition and Feed Technology
Faculty of Animal Science Bogor Agricultural University
Jalan Raya Rasamala, Bogor 16680 Indonesia
anu_jayanegara@yahoo.com*

(Makalah masuk 12 November 2012 – Diterima 19 Januari 2013)

ABSTRACT

Large amounts of polyunsaturated fatty acids undergo transformation processes in the rumen through microbial biohydrogenation to form fatty acids with higher saturation degree. The respective process explains the high content of saturated fatty acids in products of ruminants and the potential risk of consumers' health by consuming such products. Various nutritional approaches have been attempted to modulate biohydrogenation process in order to obtain healthier fatty acid profile from consumers' perspective. The present paper is aimed to review the influence of dietary tannin, a naturally produced plant secondary compound, on the pattern of polyunsaturated fatty acids biohydrogenation occurring in the rumen. The effect of tannin on some key fatty acids involved in biohydrogenation process is presented together with the underlying mechanisms, particularly from up-to-date research results. Accordingly, different form of tannin as well as different level of the application are also discussed.

Key words: Tannin, biohydrogenation, fatty acid, ruminant

ABSTRAK

POLA BIOHIDROGENASI ASAM LEMAK TAK JENUH MAJEMUK DI RUMEN DIPENGARUHI OLEH KANDUNGAN TANIN PAKAN

Sebagian besar kandungan asam lemak tak jenuh majemuk dalam pakan mengalami proses transformasi di dalam rumen melalui biohidrogenasi oleh mikroba membentuk asam lemak dengan derajat kejenuhan yang lebih tinggi. Proses biohidrogenasi inilah yang kemudian menjelaskan mengenai tingginya kandungan asam lemak jenuh dalam produk ternak ruminansia dan potensi risiko kesehatan konsumen yang mengkonsumsi produk tersebut. Berbagai pendekatan nutrisi telah dilakukan dalam rangka memodulasi proses biohidrogenasi untuk mendapatkan profil asam lemak yang menunjang kesehatan konsumen. Tulisan ini bertujuan untuk mengkaji pengaruh senyawa metabolit sekunder tanin dalam pakan terhadap pola biohidrogenasi asam lemak tak jenuh majemuk yang terjadi di dalam rumen. Pengaruh tanin pada beberapa asam lemak kunci yang terlibat dalam proses biohidrogenasi disajikan bersama dengan mekanisme yang terkait, khususnya dari hasil-hasil penelitian terkini. Terkait dengan proses biohidrogenasi tersebut, bentuk tanin yang berbeda serta pengaruh level penggunaannya atau konsentrasinya dalam pakan juga dibahas.

Kata kunci: Tanin, biohidrogenasi, asam lemak, ruminan

INTRODUCTION

Current goals of livestock production systems have expanded towards a wider context, i.e. from merely focusing on improving productivity of the animals (increasing daily weight gain, milk production, etc.) towards a more integrated approach by considering other aspects related to the systems. For instance, an important issue is regarding the role of livestock in minimizing the impact of global warming due to greenhouse gasses accumulation in the atmosphere, particularly methane emission (Cottle et al. 2011). Another current important issue that has to be

taken into account in relation to livestock production is concerning the role of livestock to promote human health through the production of animal-source foods with a particular dietetic quality. This attribute has been positively assigned to polyunsaturated fatty acids (PUFA), particularly to *n*-3 fatty acids (Barcelo-Coblijn and Murphy 2009), and conjugated linoleic acid (CLA) Benjamin and Spener (2009), but has been negatively assigned to saturated fatty acids (SFA). Unfortunately, large amounts of PUFA undergo transformation processes in the rumen through microbial biohydrogenation to form fatty acids (FA) with higher saturation degree (Chilliard et al. 2007).

The respective process explains the high content of SFA in products of ruminants and the potential risk of consumers' health by consuming such products. Therefore, the biohydrogenation process needs to be controlled in order to obtain better FA profiles in the products of animals. Accordingly, developing nutritional strategies for optimizing the respective process while maintaining optimal rumen function and fermentation is highly desirable.

The present paper is aimed to review the influence of dietary tannin, a naturally produced plant secondary compound, on the pattern of PUFA biohydrogenation occurring in the rumen. Since the respective compound is abundantly available in various tropical plants, therefore, it is of importance to be further discussed especially regarding its potential applicability to improve the quality of animal-source foods. An overview about lipid digestion in ruminants is presented in the next section, followed by some nutritional measures that have been applied to modify FA biohydrogenation in the rumen. Finally, the influence of tannin on such biohydrogenation process is presented.

LIPID DIGESTION IN RUMINANTS: LIPOLYSIS AND FATTY ACID BIOHYDROGENATION

Dietary lipids, which mainly consist of triglycerides, phospholipids and galactolipids, undergo transformation processes in the digestive tract of ruminants, i.e. through lipolysis and biohydrogenation by rumen microorganisms. Lipolysis, which is a precondition prior to the next process, results in the release of free FA, followed by biohydrogenation to saturated FA which is the reduction of the number of double bonds in the carbon chain of the FA (Jenkins 1993). The transformation processes explain the differences in the tissue profiles of FA between ruminants and non-ruminants, in which the former is more saturated than the later (Doreau et al. 2011). During lipolysis, hydrolysis of lipids is conducted mainly by microbial lipases, although there is also the action of plant lipases. The main species of rumen bacteria responsible for lipolysis is *Anaerovibrio lipolytica*. This species produces two hydrolytic enzymes, i.e. cell-bound esterase and extracellular lipase, which have activity on triglycerides and esterified fatty acids (Jenkins et al. 2008). However, *A. lipolytica* lacks the capability of hydrolyzing phospho- and galactolipids, and for these substrates, the hydrolysis is done by *Butyrivibrio* spp (Lourenco et al. 2010). In addition to bacteria, protozoal species such as *Epidinium ecaudatum* also play a considerable role in lipolysis and may contribute up to 40% of the hydrolytic activity (Lourenco et al. 2010).

After being liberated as free FA, the unsaturated FA start to enter biohydrogenation pathways. Fatty acids present in forages, cereals and oil seeds mainly contain 18 C-atoms, i.e. oleic acid (C18:1 *n*-9), linoleic acid (C18:2 *n*-6) and α -linolenic acid (C18:3 *n*-3). Biohydrogenation transforms these FA into various isomers of PUFA and monounsaturated fatty acids (MUFA), particularly *trans*-FA and CLA, and they end up as stearic acid (C18:0) which is the final product of the process (Jenkins et al. 2008). This has been demonstrated by the huge disappearance proportion of C18:2 *n*-6 and C18:3 *n*-3 in the rumen with the values of 85% and 93%, respectively (Doreau and Ferlay 1994), as well as C18:0 accumulation (Fievez et al. 2007). Among the main biohydrogenation intermediates are rumenic acid (*c*9,*t*11 C18:2), a CLA isomer, and vaccenic acid (*t*11 C18:1). The initial step of biohydrogenation is the isomerization by moving the *cis*-12 bond to either the C11 or the C13 position, followed by hydrogenation of the double bonds. Although there are several biohydrogenation pathways of C18:2 *n*-6, the main pathway is: *c*9,*c*12 C18:2 → *c*9,*t*11 C18:2 → *t*11 C18:1 → C18:0. Similarly, the main biohydrogenation pathway of C18:3 *n*-3 is: *c*9,*c*12,*c*15 C18:3 → *c*9,*t*11,*c*15 C18:3 → *t*11,*c*15 C18:2 → *t*11 C18:1 → C18:0 (Chilliard et al. 2007).

The stepwise biohydrogenation in the rumen is carried out by different groups of microorganisms, i.e. bacteria, protozoa and anaerobic fungi. *Butyrivibrio fibrisolvens* is a main bacteria species that performs biohydrogenation from C18:2 *n*-6 to *c*9,*t*11 C18:2 and *t*11 C18:1, but not to C18:0, the ultimate biohydrogenation product. From the phylogenetic tree based on 16S rRNA sequence analysis, the bacterial species that is capable of producing C18:0 from C18:2 *n*-6 is *Clostridium proteoclasticum* (Jenkins et al. 2008). Protozoa are considered to contribute significantly to FA metabolism in the rumen since about three quarters of the microbial FA originate from protozoa, considering that up to half of the rumen microbial biomass is also from protozoa. However, it has been mentioned that there was only a slight decrease in biohydrogenation after defaunation, and the presence of protozoa was not essential for the process (Lourenco et al. 2010). In the case of anaerobic fungi, especially also *Neocallimastix frontalis* is able to metabolize C18:2 *n*-6 to form *c*9, *t*11 C18:2, but the activity is very small compared to *B. fibrisolvens* (Jenkins et al. 2008).

NUTRITIONAL MEASURES FOR MANIPULATING FATTY ACID BIOHYDROGENATION IN THE RUMEN

Increasing the contents of PUFA especially of omega-3 FA in ruminant products is desirable. This is

due to, in contrast to the SFA, the presumed health benefits of omega-3 FA such as reducing the risk of cardiovascular disease in humans and lowering plasma cholesterol level (Barcelo-Coblijn dan Murphy 2009). Promoting the level of *c9, t11* C18:2, which is produced from partial biohydrogenation of C18:2 *n-6* in the rumen, is of interest since it has been shown to prevent cancer proliferation, decreased atherosclerosis and improved immune response (Palmquist et al. 2005). In relation to the respective CLA isomer, *t11* C18:1 is also desirable since it can be converted in the tissues of ruminants to *c9,t11* C18:2 via the action of the enzyme Δ -9 desaturase (stearoyl-CoA desaturase) by adding a *cis*9-double bond (Chilliard et al. 2007). Therefore, any nutritional measure for manipulating biohydrogenation of FA in the rumen is directed towards these objectives. Feeding forages to ruminants is associated with an improvement of FA profiles in the products since, in general, lipids in forages are naturally rich in PUFA (Lourenco et al. 2008). Doreau et al. (2011), for instance, mentioned that feeding grass-based diets resulted in higher levels of C18:3 *n-3* in muscle as compared to concentrate-based diets. Another study observed that feeding of grass silage to fattening bulls led to higher C18:3 *n-3* in adipose tissue than that of maize silage (Staerfl et al. 2011). Sources of variation in the rates and extents of ruminal biohydrogenation of dietary FA from forages as well as their transfer to products are forage species, cultivar, conservation method and level of inclusion (Dewhurst et al. 2006).

Supplementation of lipid rich feeds, either in the form of extracted lipids or whole oil seeds, is another strategy for modulating ruminal biohydrogenation as well as for increasing energy contents of the diets. In addition to high PUFA contents of many lipid sources (e.g., linseed, sunflower, soybean oils and seeds, fish oil), it has been known that dietary lipids may cause antimicrobial effects on biohydrogenating bacteria especially lipids with higher degree of unsaturation. Lipids rich in either C18:2 *n-6* or C18:3 *n-3* are both contributing to the increase of CLA contents in milk (Lourenco et al. 2010). Fish oil, which is rich in long-chain PUFA, particularly eicosapentaenoic acid (EPA, C20:5 *n-3*) and docosahexaenoic acid (DHA, C22:6 *n-3*), has been shown to inhibit the last step of biohydrogenation, i.e. the conversion of *t11* C18:1 to C18:0, which may be due to its influence on *B. fibrisolvans*, *Ruminococcus albus* and *Selenomonas ruminantium* (Potu et al. 2011). Alternatively to fish oil, microalgae also contain long-chain PUFA and its supplementation increased C22:6 *n-3*, CLA and *t11* C18:1 levels in milk (Boeckaert et al. 2008). Several oils that are rich in MCFA such as coconut (rich in lauric acid, C12:0, and myristic acid, C14:0) and palm oils (rich in palmitic acid, C16:0) are considered not to

have benefited FA composition in ruminant products (Lourenco et al. 2010). Administration of lipids in the form of whole seeds usually (but not always) results in a lower biohydrogenation than when given as extracted oils. The extent of this reduction depends on the hardness of the hull, particle size and mastication for breaking the oilseed hulls (Doreau et al. 2011). Prior to supplementation, lipids may also be protected through various treatments. The ideal criteria of lipid protection technology have been proposed by Jenkins and Bridges Jr. (2007) as (1) consistent and predictable enhancement of UFA flow to the duodenum, (2) adequate release and absorption of the UFA in the intestines, and (3) minimal adverse effects on ruminal fermentation. Some protection techniques that have been implemented are, for instance, encapsulation of emulsified oils into a matrix of formaldehyde-treated proteins (Sterk et al. 2010), formation of calcium salts (Theurer et al. 2009) and formation of fatty acyl amides (Lundy et al. 2004).

Plant secondary compounds may alter ruminal biohydrogenation of FA as well. Studies of Leiber et al. (2004 and 2005) showed that C18:3 *n-3* levels in milk were to some extent independent of their intake. Accordingly, the results then led to a hypothesis that plant secondary compounds may play a role to the elevated contents of C18:3 *n-3* and/or *c9,t11* C18:2 observed in milk and milk products originating from cows grazing on alpine pastures compared to those from intensive lowland production systems. These compounds may include polyphenol or tannin (will be discussed in the next section), saponin and essential oil. Regarding saponin, it has been reported that a saponin-rich extract from *Yucca schidigera* inhibited the growth of *B. fibrisolvans* more than other bacteria species (Wallace et al. 1994), suggesting that there is a potential of saponins in modifying FA biohydrogenation. However, further research needs to be undertaken to ascertain this since direct evidence so far has shown lack of effects of saponins in altering FA composition in ruminal fluid (Khiaosa-Ard et al. 2009), milk (Benchaar and Chouinard 2009) and meat (Brognna et al. 2011). Essential oils have an activity against several bacteria that are involved in ruminal biohydrogenation. Their effects in the rumen are quite variable, depending on nature of the compounds, level of applications, basal diet and adaptation time of ruminal microbes in the presence of the compounds (Benchaar et al. 2008). Some plant extracts and essential oils from Australian plants have been reported to have a selective inhibitory effect on *C. proteoclasticum* without affecting *B. fibrisolvans*, and some can inhibit the saturation of C18:2 *n-6*, *c9,t11* C18:2 and *t11* C18:1 *in vitro* (Durmic et al. 2008). On the contrary, feeding of cinnamaldehyde, the main component of cinnamon bark (*Cinnamomum cassia*)

essential oil to dairy cows revealed no effects on milk fatty acid profile thus suggesting that the potential of this essential oil to alter ruminal biohydrogenation process is low (Benchaar and Chouinard 2009).

TANNIN AND ITS INFLUENCE ON FATTY ACID BIOHYDROGENATION

Tannin can be divided into two categories based on its property in binding protein, i.e. tannin phenol (or tannin) which can bind protein, and non-tannin phenol which is not able to bind protein. Generally, tannin has a much higher molecular weight than that of non-tannin phenol; the latter is simpler phenol such as catechol, pyrogallol, gallic acid, catechin and other flavanols (Scalbert 1991). Based on its molecular structure, tannin is classified as hydrolysable tannin (HT) and condensed tannin (CT; also known as proanthocyanidin), although other tannin may occur which is combination of these two basic structures (McSweeney et al. 2001). There is a great structural diversity of tannin between different sources and even slight changes in the structure may produce measurable biological effects (Mueller-Harvey 2006). Hydrolysable tannin contains carbohydrate (generally glucose molecule) as a central core with hydroxyl groups which are esterified with phenolic groups. Different from that, CT does not have a central carbohydrate core and are complexes of oligomers and polymers of flavanoid units linked by carbon-carbon bonds with a molecular weight of 2,000-4,000 kDa (Goel et al. 2005). Hydrolysable tannin is more susceptible to enzymatic and non-enzymatic hydrolysis than that of CT, and usually is better soluble in water. Further, based on the hydrolysis products, HT can be divided into gallotannin, which yield gallic acid and glucose, and ellagitannin, which yield ellagic acid and glucose (Reed 1995).

Multiple phenolic hydroxyl groups enable tannin to form complexes especially with protein, although complexes with polysaccharides and minerals are also occurring. With regard to interactions between tannin and protein, the strengths of the complexes depend on characteristics of both tannin and protein such as molecular weight, tertiary structure, isoelectric point and compatibility of binding sites (Reed 1995). Three different bonds may exist in tannin-protein complexes, i.e. hydrogen bonds through a large number of free phenolic hydroxyl groups, hydrophobic bonds through aromatic ring structures and covalent bonds through oxidative polymerization reactions due to heating, exposure to UV radiation and the action of polyphenol oxidase (Silanikove et al. 2001).

In relation to biohydrogenation of FA in the rumen, tannin has been reported to reduce PUFA

biohydrogenation, accumulate t_{11} C18:1 and/or decrease concentration of C18:0 (Khiaosa-Ard et al. 2009; Vasta et al. 2009; Cabiddu et al. 2010; Jayanegara et al. 2011; Jayanegara et al. 2012), to increase the transfer rate of C18:3 $n-3$ from feed to milk as a product (Kälber et al. 2011), and thus may alter FA composition in the products of animals (Doreau et al. 2011). *In vitro* incubations of tropical (Jayanegara et al. 2011) and alpine plants (Jayanegara et al. 2012) with additional linseed oil suggest the ability of plant phenolics in modulating FA biohydrogenation, i.e. by decelerating the process right from the first step. This was indicated by lower disappearance of C18:3 $n-3$ and C18:2 $n-6$ in the incubations of both tropical and alpine plants containing high phenolics in the respective studies. Such effects may be explained by the toxicity of phenolics, particularly of tannin phenolics, on bacterial species involved in FA biohydrogenation (Khiaosa-Ard et al. 2009; Vasta et al. 2010). It is also possible that phenolics may inhibit the process of lipolysis which is a precondition prior to further transformation of fatty acids, i.e. biohydrogenation (Jenkins 1993; Lourenco et al. 2010). In accordance to this, Cabiddu et al. (2010) observed a negative relationship between tannins present in *Vicia sativa* and *Trifolium incarnatum* with the biohydrogenation of C18:3 $n-3$. Other phenolics considered to be responsible for the increase found in the transfer of C18:3 $n-3$ from feed to milk (Kälber et al. 2011) when fed buckwheat, which might be rutin or fagopyrin (Leiber et al. 2012).

Regarding the occurrence of biohydrogenation intermediates by the influence of plant phenolics, in an *in vitro* study, Khiaosa-Ard et al. (2009) observed a considerable increase of t_{11} C18:1 at the expense of C18:0 when adding *Acacia mearnsii* extract (source of CT) at 79 g/kg DM to a grass-clover hay diet supplemented with linseed oil. The authors therefore suggested that the tannins inhibited the terminal step of FA biohydrogenation. The accumulation of t_{11} C18:1 in the presence of tannin phenolics was confirmed by other *in vitro* study (Vasta et al. 2009). This effect might be related to phenolic toxicity, especially of tannin, on bacterial species involved in FA biohydrogenation, through selective inhibition of cell wall synthesis (Smith et al. 2005), interaction of phenols with microbial proteins (Silanikove et al. 2001), and direct interaction between phenols and lipids (He et al. 2006). *Clostridium proteoclasticum*, a bacterial species responsible for the terminal step of biohydrogenation, i.e. the conversion from t_{11} C18:1 to C18:0 (Jenkins et al. 2008), was proved to decrease by 31% in the rumen of lambs fed by a tannin-supplemented diet compared to a control diet (Vasta et al. 2010).

Different fractions of tannin, i.e. HT and CT, contributed to the modification of biohydrogenation pattern. Jayanegara et al. (2011) found that the main class of polyphenol which prevented C18:3 *n*-3 and C18:2 *n*-6 from biohydrogenation was that of the HT, while the appearance of *c*9, *t*11 C18:2 was closer correlated with the CT. This illustrates that both types of tannins are involved in the inhibition of biohydrogenation but in different steps, namely HT in the first and CT in the second step. By contrast, it has been observed *in vitro* and *in vivo* that the addition of CT extracts to the diet caused no difference in the concentration of conjugated C18:2 in the ruminal fluid, but led to a considerable increase of *t*11 C18:1 at the expense of C18:0 instead (Khiaosa-Ard et al. 2009). This indicates that inhibition of the third step of biohydrogenation took place. Thus, a differentiated and still not always coherent influence of different tannins on the biohydrogenation pathway can be observed.

It has to be noted that the effect of tannin on FA biohydrogenation cannot be found in every study (Abbeddou et al. 2011) and it appears that the phenolic effects are minor when the concentration is below a certain threshold (Jayanegara et al. 2011; Khiaosa-Ard et al. 2011). For instance, supplementation of commercially available extracts of tannins, i.e. a 1:1 mixture of *Castanea sativa* and *Schinopsis lorentzii* (quebracho, a source of CT) at a level of 10 mg/g DM to a diet containing sunflower oil did not alter the proportions of the major FA classes in milk (PUFA, MUFA and SFA) as well as the proportions of *c*9, *t*11 C18:2 and *t*11 C18:1 in milk (Toral et al. 2011). The low dose of the tannin mixture in that study was presumed to have been the reason for the lack of change in the FA in the respective study. It appears that the levels of polyphenols in diets as well as the sources from where they were obtained are among the factors of influence affecting the biohydrogenation of FA and general rumen fermentation (Jayanegara et al. 2011).

CONCLUSION

The pattern of PUFA biohydrogenation can be modulated in the presence of dietary tannin and, thus, may potentially modify FA composition in the products of animals. Among the main effects of tannin on biohydrogenation are reducing PUFA biohydrogenation, accumulating *t*11 C18:1 and/or decreasing concentration of C18:0. There is still a controversy whether the respective substance is able to increase *c*9,*t*11 C18:2 concentration in rumen fluid. The effect of tannin on biohydrogenation appears to be source and dose-dependent.

REFERENCES

- Abbeddou S, Rischkowsky B, Richter EK, Hess HD, Kreuzer M. 2011. Modification of milk fatty acid composition by feeding forages and agro-industrial byproducts from dry areas to Awassi sheep. *J Dairy Sci.* 94:4657-4668.
- Barcelo-Coblijn G, Murphy EJ. 2009. Alpha-linolenic acid and its conversion to longer chain *n*-3 fatty acids: benefits for human health and a role in maintaining tissue *n*-3 fatty acid levels. *Prog Lipid Res.* 48:355-74.
- Benchaar C, Calsamiglia S, Chaves AV, Fraser GR, Colombatto D, McAllister TA, Beauchemin KA. 2008. A review of plant-derived essential oils in ruminant nutrition and production. *Anim Feed Sci Technol.* 145:209-228.
- Benchaar C, Chouinard PY. 2009. Assessment of the potential of cinnamaldehyde, condensed tannins, and saponins to modify milk fatty acid composition of dairy cows. *J Dairy Sci.* 92:3392-3396.
- Benjamin S, Spener F. 2009. Conjugated linoleic acids as functional foods: an insight into their health benefits. *Nutr Metab.* 6:36.
- Boeckeaert C, Vlaeminck B, Fievez V, Maignien L, Dijkstra J, Boon N. 2008. Accumulation of *trans* C18:1 fatty acids in the rumen after dietary algal supplementation is associated with changes in the *Butyrivibrio* community. *Appl Environ Microbiol.* 74:6923-6930.
- Brogna DMR, Nasri S, Ben Salem H, Mele M, Serra A, Bella M, Priolo A, Makkar HPS, Vasta V. 2011. Effect of dietary saponins from *Quillaja saponaria* L. on fatty acid composition and cholesterol content in muscle *Longissimus dorsi* of lambs. *Animal* 5:1124-1130.
- Cabiddu A, Salis L, Tweed JKS, Molle G, Decandia M, Lee MRF. 2010. The influence of plant polyphenols on lipolysis and biohydrogenation in dried forages at different phenological stages: *in vitro* study. *J Sci Food Agric.* 90:829-835.
- Chilliard Y, Glasser F, Ferlay A, Bernard L, Rouel J, Doreau M. 2007. Diet, rumen biohydrogenation and nutritional quality of cow and goat milk fat. *Eur J Lipid Sci Technol.* 109:828-855.
- Cottle DJ, Nolan JV, Wiedemann SG. 2011. Ruminant enteric methane mitigation: a review. *Anim Prod Sci.* 51:491-514.
- Dewhurst RJ, Shingfield KJ, Lee MRF, Scollan ND. 2006. Increasing the concentrations of beneficial polyunsaturated fatty acids in milk produced by dairy cows in high-forage systems. *Anim Feed Sci Technol.* 131:168-206.
- Doreau M, Ferlay A. 1994. Digestion and utilisation of fatty acids by ruminants. *Anim. Feed Sci Technol.* 45:379-396.
- Doreau M, Bauchart D, Chilliard Y. 2011. Enhancing fatty acid composition of milk and meat through animal feeding. *Anim Prod Sci.* 51:19-29.

- Durmic Z, Mcsweeney CS, Kemp GW, Hutton P, Wallace RJ, Vercoe PE. 2008. Australian plants with potential to inhibit bacteria and processes involved in ruminal biohydrogenation of fatty acids. *Anim Feed Sci Technol*. 145:271-284.
- Fievez V, Vlaeminck B, Jenkins T, Enjalbert F, Doreau M. 2007. Assessing rumen biohydrogenation and its manipulation *in vivo*, *in vitro* and *in situ*. *Eur J Lipid Sci Technol*. 109:740-756.
- Goel G, Puniya AK, Aguilar CN, Singh K. 2005. Interaction of gut microflora with tannins in feeds. *Naturwissenschaften* 92:497-503.
- He Q, Shi B, Yao K. 2006. Interactions of gallotannins with proteins, amino acids, phospholipids and sugars. *Food Chem*. 95:250-254.
- Jayanegara A, Kreuzer M, Wina E, Leiber F. 2011. Significance of phenolic compounds in tropical forages for the ruminal bypass of polyunsaturated fatty acids and the appearance of biohydrogenation intermediates as examined *in vitro*. *Anim Prod Sci*. 51:1127-1136.
- Jayanegara A, Kreuzer M, Leiber F. 2012. Ruminant disappearance of polyunsaturated fatty acids and appearance of biohydrogenation products when incubating linseed oil with alpine forage plant species *in vitro*. *Livest Sci*. 147:104-112.
- Jenkins TC 1993. Lipid metabolism in the rumen. *J Dairy Sci*. 76:3851-3863.
- Jenkins TC, Bridges Jr WC. 2007. Protection of fatty acids against ruminal biohydrogenation in cattle. *Eur J Lipid Sci Technol*. 109:778-789.
- Jenkins TC, Wallace RJ, Moate PJ, Mosley EE. 2008. Recent advances in biohydrogenation of unsaturated fatty acids within the rumen microbial ecosystem. *J Anim Sci*. 86:397-412.
- Kälber T, Meier JS, Kreuzer M, Leiber F. 2011. Flowering catch crops used as forage plants for dairy cows: influence on fatty acids and tocopherols in milk. *J Dairy Sci*. 94:1477-1489.
- Khiaosa-Ard R, Bryner SF, Scheeder MRL, Wettstein HR, Leiber F, Kreuzer M, Soliva CR. 2009. Evidence for the inhibition of the terminal step of ruminal α -linolenic acid biohydrogenation by condensed tannins. *J Dairy Sci*. 92:177-188.
- Khiaosa-Ard R, Soliva CR, Kreuzer M, Leiber F. 2011. Influence of alpine forage either employed as donor cow's feed of ras incubation substrate on *in vitro* ruminal fatty acid biohydrogenation. *Livest Sci*. 140:80-87.
- Leiber F, Kreuzer M, Jörg B, Leuenberger H, Wettstein HR. 2004. Contribution of altitude and Alpine origin of forage to the influence of Alpine sojourn of cows on intake, nitrogen conversion, metabolic stress and milk synthesis. *Anim Sci*. 78:451-466.
- Leiber F, Kreuzer M, Nigg D, Wettstein HR, Scheeder MRL. 2005. A study on the causes for the elevated *n*-3 fatty acids in cows' milk of Alpine origin. *Lipids* 40:191-202.
- Leiber F, Kunz C, Kreuzer M. 2012. Influence of different morphological parts of buckwheat (*Fagopyrum esculentum*) and its major secondary metabolite rutin on rumen fermentation *in vitro*. *Czech J Anim Sci*. 57:10-18.
- Lourenco M, Van Ranst G, Vlaeminck B, De Smet S, Fievez V. 2008. Influence of different dietary forages on the fatty acid composition of rumen digesta as well as ruminant meat and milk. *Anim. Feed Sci Technol*. 145:418-437.
- Lourenco M, Ramos-Morales E, Wallace RJ. 2010. The role of microbes in rumen lipolysis and biohydrogenation and their manipulation. *Animal* 4:1008-1023.
- Lundy FP, Block E, Bridges WC, Bertrand JA, Jenkins TC. 2004. Ruminant biohydrogenation in Holstein cows fed soybean fatty acids as amides or calcium salts. *J Dairy Sci*. 87:1038-1046.
- Mcsweeney CS, Palmer B, McNeill DM, Krause DO. 2001. Microbial interactions with tannins: nutritional consequences for ruminants. *Anim Feed Sci Technol*. 91:83-93.
- Mueller-Harvey I. 2006. Unravelling the conundrum of tannins in animal nutrition and health. *J Sci Food Agric*. 86:2010-2037.
- Palmquist DL, Lock AL, Shingfield KJ, Bauman DE. 2005. Biosynthesis of conjugated linoleic acid in ruminants and humans. *Adv Food Nutr Res*. 50:179-217.
- Potu RB, Abu Ghazaleh AA, Hastings D, Jones K, Ibrahim SA. 2011. The effect of lipid supplements on ruminal bacteria in continuous culture fermenters varies with the fatty acid composition. *J Microbiol*. 49:216-223.
- Reed JD. 1995. Nutritional toxicology of tannins and related polyphenols in forage legumes. *J Anim Sci*. 73. 1516-1528.
- Scalbert A. 1991. Antimicrobial properties of tannins. *Phytochem*. 30:3875-3883.
- Silanikove N, Perevolotsky A, Provenza FD. 2001. Use of tannin-binding chemicals to assay for tannins and their negative postingestive effects in ruminants. *Anim. Feed Sci Technol*. 91:69-81.
- Smith AH, Zoetendal E, Mackie RI. 2005. Bacterial mechanisms to overcome inhibitory effects of dietary tannins. *Microb Ecol*. 50. 197-205.
- Staerfl SM, Soliva CR, Leiber F, Kreuzer M. 2011. Fatty acid profile and oxidative stability of the perirenal fat of bulls fattened on grass silage and maize silage supplemented with tannins, garlic, maca and lupines. *Meat Sci*. 89:98-104.
- Sterk A, Hovenier R, Vlaeminck B, Van Vuuren AM, Hendriks WH, Dijkstra J. 2010. Effects of chemically or technologically treated linseed products and docosahexaenoic acid addition to linseed oil on biohydrogenation of C18:3 *n*-3 *in vitro*. *J Dairy Sci*. 93:5286-5299.

- Theurer ML, Block E, Sanchez WK, Mcguire MA. 2009. Calcium salts of polyunsaturated fatty acids deliver more essential fatty acids to the lactating dairy cow. *J Dairy Sci.* 92:2051-2056.
- Toral PG, Hervas G, Bichi E, Belenguer A, Frutos P. 2011. Tannins as feed additives to modulate ruminal biohydrogenation: effects on animal performance, milk fatty acid composition and ruminal fermentation in dairy ewes fed a diet containing sunflower oil. *Anim. Feed Sci Technol.* 164:199-206.
- Vasta V, Makkar HPS, Mele M, Priolo A. 2009. Ruminal biohydrogenation as affected by tannins *in vitro*. *Brit J Nutr.* 102:82-92.
- Vasta V, Yanez-Ruiz DR, Mele M, Serra A, Luciano G, Lanza M, Biondi L, Priolo A. 2010. Bacterial and protozoal communities and fatty acid profile in the rumen of sheep fed a diet containing added tannins. *Appl Environ Microbiol.* 76:2549-2555.
- Wallace RJ, Arthaud L, Newbold CJ. 1994. Influence of *Yucca schidigera* extract on ruminal ammonia concentrations and ruminal microorganisms. *Appl Environ Microbiol.* 60:1762-1767.